arXiv:1610.03046v2 [math.GT] 29 Dec 2016

The classification of Kleinian groups of Hausdorff dimensions at most one

Yong Hou*

Institute for Advanced Study

Abstract

In this paper we provide the complete classification of Kleinian groups of Hausdorff dimensions less than 1. In particular, we prove that every purely loxodromic Kleinian groups of Hausdorff dimension < 1 is a classical Schottky group. This upper bound is sharp. As an application, the result of [5] then implies that, every closed Riemann surface is uniformizable by a classical Schottky group. The prove relie on the result of Hou [7], and space of rectifiable Γ-invariant closed curves.

1 Introduction and Main Theorem

We take Kleinian groups to be finitely generated, torsion-free, discrete subgroups of $\mathrm{PSL}(2,\mathbb{C})$. The main theorem is:

Theorem 1.1 (Classification)**.** *Any purely loxodromic Kleinian group Γ with limit set of Hausdorff dimension < 1 is a classical Schottky group. This bound is sharp.*

We note that by Selberg lemma, torsion-free is not really a restriction, since any finitely generated discrete subgroup of $\mathrm{PSL}(2,\mathbb{C})$ have finite index torsion-free subgroup.

*Supported by Ambore Mondell Fundation

As an application, we have the following Corollary 1.3, which is an resolution of a folklore problem of Bers on classical Schottky group uniformization of closed Riemann surface. Corollary 1.3 follows from the work of Hou [5]:

Theorem 1.2 (Hou [5])**.** *Every closed Riemann surface is uniformizable by a Schottky group of Hausdorff dimension* < 1*, i.e. every point in moduli space has a Hausdorff dimension* < 1 *fiber in the Schottky space.*

Corollary 1.3. *(Uniformization) Every closed Riemann surface can be uniformized by a classical Schottky group.*

1.1 Strategy of proof

First let us recall the result of Hou [7]:

Theorem 1.4 (Hou)**.** *There exists* $\lambda > 0$ *such that any Kleinian group with limit set of Hausdorff dimension* $< \lambda$ *is a classical Schottky group*

Define $H_c = \sup\{\lambda | \text{satisfies Theorem 1.4}\}$. H_c is the maximal parameter such that if Γ have Hausdorff dimension $< H_c$ then Γ is classical Schottky group. Hence Theorem 1.1 can be rephrased as:

Remark 1.5. $H_c \geq 1$.

We prove by contradiction, assume that $H_c < 1$.

First, it is consequence of Theorem 1.4 that, $\mathfrak{J}_{g,o}^{H_c}$ (rank-g classical Schottky groups of Hausdorff dimension $< H_c$, see Section 2) is a $3g-3$ dimensional complex open submanifold of $\mathfrak{J}_g$, the rank-g Schottky space. By definition of H_c, there must exist Γ non-classical Schottky group, in the boundary $\partial\mathfrak{J}_{g_o}^{H_c}$, of Hausdorff dimension H_c. Hence, we have a sequence of classical Schottky groups $\{\Gamma_n\} \subset \mathfrak{J}_{g,o}^{H_c}$ such that, $\Gamma_n \to \Gamma$ in Schottky space $\mathfrak{J}_g$.

Next, it is a result of Bowen [2] that, a Schottky group Γ of Hausdorff dimension < 1, if and only if there exist a *rectifiable* Γ-invariant closed curve. Let $\mathscr{R}(S^1, W)$ be the space of rectifiable closed curves in compact set $W \subset \mathbb{C}$ with Frechet metric, see Section 2. This is complete metric space [1]. We show that if $H_c < 1$ then, every Γ-invariant closed curve is limit of Γ_n-invariant closed curves in $\mathscr{R}(S^1, W)$. And we also show that if Γ is also non-classical Schottky group, then there exists a sequence of rectifiable Γ_n-invariant closed curves that do not converge to any Γ-invariant closed curve of Γ.

This non-limit point is proved by constructing a sequence of Γ_n-invariant closed curves which are called *linear* that rely on the the property of Γ_n being classical Schottky groups. When Γ is non-classical schottky group, we use the crucial fact that, every generating set of Γ must have at least a non-circle for the generating Jordan curves. Since all generating sets are equivalent under Nielsen transformations, this property of "wiggling of some Jordan curve" will persist to all subsequent level of transformations, which will preclude eventual Γ-invariant closed curve to be limit of linear invariant closed curves.

Acknowledgement

This work is made possible by unwavering supports and brilliantly insightful conversations from Peter Sarnak, whom I'm greatly indebted to. It is the groundbreaking works of Peter Sarnak that has guided the author to study this problem at first. I'm very grateful to the advices and, guides and, ideas that Peter provided for this work. I wish to express my deep gratitude and sincere appreciation to Dave Gabai, Benson Farb, and Ian Agol for continuous amazing support and many fundamentally influential conversations which greatly impacted my works.

I want to express my greatest appreciation to Ian Agol for very detailed reading of the previous draft and help me corrected many mistakes that includes many notational and statement issues. In addition, my thanks to Mary Yan He for helpful comments on the paper.

This work is dedicated to my parents: Shu Ying Hou and Sulan Sun.

2 Generating Jordan curves and $\mathscr{R}(S^1, W)$

Schottky group Γ of rank g is defined as convex-cocompact discrete faithful representation of $\mathbb{F}_g$ in $\mathrm{PSL}(2,\mathbb{C})$. It follows that Γ is freely generated by purely loxodromics $\{\gamma_i\}_1^g$. This implies we can find collection of open topological disks $D_i, D_{i+g}, 1 \leq i \leq g$ of disjoint closure $\bar{D}_i \cap \bar{D}_{i+g} = \emptyset$ in the Riemann sphere $\partial\mathbb{H}^3 = \overline{\mathbb{C}}$ with boundary curves $\partial\bar{D}_i = c_i, \partial\bar{D}_{i+g} = c_{i+g}$. By definition c_i, c_{i+g} are closed Jordan curves in Riemann sphere $\partial\mathbb{H}^3$, such that $\gamma_i(c_i) = c_{i+g}$ and $\gamma_i(D_i) \cap D_{i+g} = \emptyset$. Whenever there exists a set $\{\gamma_1, ..., \gamma_g\}$ of generators with all $\{c_i, c_{i+g}\}_1^g$ as circles, then it is called a classical Schottky group with $\{\gamma_1, ..., \gamma_g\}$ classical generators.

Schottky space $\mathfrak{J}_g$ is defined as space of all rank g Schottky groups up to conjugacy by $\mathrm{PSL}(2,\mathbb{C})$. By normalization, we can chart $\mathfrak{J}_g$ by $3g-3$ complex parameters. Hence $\mathfrak{J}_g$ is $3g-3$ dimensional complex manifold. The

bihomolomorphic Auto($\mathfrak{J}_g$) group is Out($\mathbb{F}_g$), which is isomorphic to quotient of the handle-body group. Denote by $\mathfrak{J}_{g,o}$ the set of all elements of $\mathfrak{J}_g$ that are classical Schottky groups. On the other hand, $\mathfrak{J}_{g,o}$ is not submanifold. In fact, it is nontrivial result due to Marden that $\mathfrak{J}_{g,o}$ is non-empty and non-dense subset of $\mathfrak{J}_g$. However, it follows from Theorem 1.4, $\mathfrak{J}^{\lambda}_{g,o}$ is $3g-3$ dimensional complex submanifold. Here $\mathfrak{J}^{\lambda}_{g,o}$ denotes space of classical Schottky groups of Hausdorff dimension $< \lambda$.

Let Γ be a Schottky group generated by $S = \{\gamma_1, ..., \gamma_g\}$. We define generating Jordan curves space $\mathscr{C}_S \subset \{h : S^1 \to W\}$, for h continuous maps, to be the collection of all C_g Jordan closed curves $C_g = \{c_1, ..., c_{2g}\}$ such that pair-wisely $\gamma_i(c_i) = c_{i+g}$ and $\gamma_i(D_i) \cap D_{i+g} = \emptyset$, for $1 \leq i \leq g$ where D_i is the topological open disk bounded by c_i and disjoint from c_j, for all $j \neq i$.

Let $\mathscr{D}$ denote the space of all continuous functions f on $\overline{\mathbb{C}}$ which is harmonic on $\mathbb{C}\setminus c_i$, and with finite positive Dirichlet integral on $\mathbb{C}\setminus c_i$. For $z \in \mathbb{C}$, $z = x + \sqrt{-1}y$ the Dirichlet integral $\Delta^{\pm}(f, c_i)$ of f on $\mathbb{C} \setminus c_i$ is defined as:

$$\Delta^{-}(f, c_i) = \int\int_{D_i} (\partial_x f)^2 + (\partial_y f)^2 \mathrm{d}x\mathrm{d}y - \int\int_{\overline{D}_i^c} (\partial_x f)^2 + (\partial_y f)^2 \mathrm{d}x\mathrm{d}y,$$

$$\Delta^{+}(f, c_i) = \int\int_{D_i} (\partial_x f)^2 + (\partial_y f)^2 \mathrm{d}x\mathrm{d}y + \int\int_{\overline{D}_i^c} (\partial_x f)^2 + (\partial_y f)^2 \mathrm{d}x\mathrm{d}y.$$

Let $\sigma_{c_i} = \sup_{\mathscr{D}} \frac{|\Delta^{-}(f,c_i)|}{\Delta^{+}(f,c_i)}$. It follows from [8], σ_{c_i} is Mobius invariant and $\sigma_{c_i} \in [0, 1]$, such that $\sigma_{c_i} = 0$ if and only if c_i is a circle.

Define a map $\psi : \mathscr{C}_S \to [0, g]$ given by $\psi(C_g) = \sum_{i=1}^{g} \sigma_{c_i}$. Set $\theta_S = \inf_{\mathscr{C}_S} \psi(C_g)$, and $\mathscr{C}^*_S = \psi^{-1}(\theta_S)$. Note that if $\theta_S > 0$ then S is not classical generating set of Γ.

Note that one should be careful that it is possible that $\mathscr{C}^*_S$ is empty when Γ is a nonclassical Schottky group generated by tangent circles. However this won't effect us since we will only be using $\mathscr{C}^*_S$ when Γ is a classical Schottky group.

Notation:

- In general we always arrange elements of C_g so that c_i, c_{i+g} are the corresponding generating Jordan curves of generator γ_i. However, sometimes for convenience we will denote a pair of Jordan generating curves of C_g corresponding to a generator γ by subscribes $\pm$ such as: $c_{\gamma,-}, c_{\gamma,+}$ or with generator omitted as c_-, c_+. For $\gamma \in S$, we also set $\sigma_\gamma = \sigma_{c_{\gamma,-}}$.

- Given Γ a Kleinian group, we denote by Λ_Γ and Ω_Γ and $\mathfrak{D}_\Gamma$ its limit set, region of discontinuity, and Hausdorff dimension respectively throughout this paper.

Definition 2.1. Given a Schottky group Γ, a closed Γ-invariant Jordan curve that contains the limit set Λ_Γ is called *quasi-circle* of Γ.

Remark: From now on, we make the global assumption throughout this paper that *all classical Schottky groups* are of Hausdorff dimension $\mathfrak{D}_\Gamma < H_c$, if not stated otherwise.

Next we give a construction of quasi-circles of Γ which is a generalization of the construction by Bowen [2].

For $C_g \in \mathscr{C}_S$, let $a_{p,q}$ denote collection of arcs $\{a_{i,p_i,q_i}\}$ connecting points $p_i \in c_i, q_i \in c_{i+g}, c_i, c_{i+g} \in C_g$ for $1 \leq i \leq g$, and arc on c_{i+g} that connects q_i to $\gamma_i(p_i)$. So $a_{p,q}$ is a set of g disjoint curves connecting disjoint points on collection of Jordan curves of C_g.

$\eta_\Gamma(a_{p,q}) = \Lambda_\Gamma \cup \cup_{\gamma\in\Gamma}\gamma(a_{p,q})$ defines a Γ-invariant closed curve containing Λ_Γ. $\eta_\Gamma(a_{p,q})$ defines a quasi-circle of Γ. Of course, $\eta_\Gamma(a_{p,q})$ are not unique and different $a_{p,q}$ gives distinct quasi-circles. We call $\eta_\Gamma(a_{p,q})$ a *quasi-circle with marking* C_g. The simply connected regions $\mathbb{C} \setminus \eta_\Gamma(a_{p,q})$ gives Bers simultaneous uniformization of Riemann surface Ω_Γ/Γ.

Note that the quasi-circles constructed in [2], which requires that q_i is a imagine of p_i under element of $\gamma_i \in S$, is a subset of the collection that we have defined here. In fact, this generalization is crucial for the construction of quasi-circles of non-classical Schottky groups. In addition, we also have the following.

Proposition 2.2. *Every quasi-circle of* Γ *can be written as* $\eta_\Gamma(a_{p,q})$ *for some marking and* $a_{p,q}$.

Proof. Let η_Γ be a quasi-circle of Γ. Since η_Γ contains Λ_Γ, let $\rho_\Gamma = \eta_\Gamma \setminus \Lambda_\Gamma$. Since η_Γ is Γ-invariant, we have ρ_Γ is Γ-invariant disjoint curves. ρ_Γ is open and $\rho_\Gamma \subset \Omega_\Gamma$, let F_Γ be a fundamental domain of Ω_Γ, then $\sigma_\Gamma = \rho_\Gamma \cap F_\Gamma$ is a fundamental domain of ρ_Γ such that, $\Gamma(\sigma_\Gamma) = \rho_\Gamma$.

Since F_Γ of Γ are given by some generating set S, hence defines $C_g \in \mathscr{C}_S$. Let $b = \cup_i^g(c_i \cup c_{i+g}) \cap \sigma_\Gamma$ and $a = \sigma_\Gamma \setminus b$. Since $a = \{p_i, q_{i+g}\}_{1\leq i\leq g}$ for $p_i, q_{i+g} \in b$ hence it defines a disjoint curve collection of η_Γ. Therefore, σ_Γ is the desired generating curve collection for η_Γ. □

Definition 2.3. For a given marked quasi-circle $\eta_\Gamma(a_{p,q})$ we define the j-level $\eta^j_{a,p}$ of $\eta_\Gamma(a_{p,q})$ as the curve segments of $\eta_\Gamma(a_{p,q})$ which are images of $a_{p,q}$ under word of S of length j with respect to the C_g generating curves.

Definition 2.4. We call a quasi-circle $\eta_\Gamma(a_{p,q})$ *linear* if either it is piece-wise circular arc or piece-wise line. ln particular if $\eta^j_\Gamma(a_{p,q})$ are piece-wise circular arc or lines.

In general we denote quasi-circles of Γ by η_Γ. We use $\eta_\Gamma(a_{p,q})$ to denote quasi-circles constructed by inital curves $a_{p,q}$ as defined earlier.

It should be noted that a quasi-circle of Γ in general is not necessarily rectifiable. For instance, if we take $a_{p,q}$ to be some non-rectifiable curves then, $\eta_\Gamma(a_{p,q})$ will be non-rectifiable. This is not the only obstruction to rectifiability, in fact we have the following result:

Theorem 2.5 ([2])**.** *For a given Schottky group* Γ*, the Hausdorff dimension of limit set is* $\mathfrak{D}_\Gamma < 1$*, if and only if there exists a rectifiable quasi-circle for* Γ*.*

The proof of Theorem 2.5 relies on the fact that the Poincare series of Γ converges if and only if $\mathfrak{D}_\Gamma < 1$ [2].

Proposition 2.6. *Let* Γ *be a Schottky group of* $\mathfrak{D}_\Gamma < 1$*. Suppose* $a_{p,q}$ *is a rectifiable curve. Then* $\eta_\Gamma(a_{p,q})$ *is a rectifiable quasi-circle of* Γ*.*

Proof. Let μ^1 be the 1-dimensional Hausdorff measure. Since $\mathfrak{D}_\Gamma < 1$, we have $\mu^1(\Lambda_\Gamma) = 0$. Let $\Gamma = \{\gamma_k\}_{k=1}^\infty$ we have,

$$\begin{aligned}\mu^1(\eta_\Gamma(a_{p,q})) &= \sum_{k=1}^\infty \mu^1(\gamma_k(a_{p,q})) \\ &\asymp \mu^1(a_{p,q}) \sum_{k=1}^\infty |\gamma_k'(z)|, \quad z \in a_{p,q}.\end{aligned}$$

Also by $\mathfrak{D}_\Gamma < 1$, the Poincare series satisfies $\sum_{k=1}^\infty |\gamma_k'| < \infty$. This implies that $\eta_\Gamma(a_{p,q})$ is rectifiable if and only if $a_{p,q}$ rectifiable. □

Let $W \subset \mathbb{C}$ be a compact set. Denote the space of closed *rectifiable* curves in $\mathbb{C}$ that intersect with W by

$$\mathscr{R}(S^1, W) \subset \{h : S^1 \to \mathbb{C} | h(S^1) \cap W \neq \emptyset, \quad h \text{ continuous rectifiable map.}\}$$

For $h_1, h_2 \in \mathscr{R}(S^1, W)$ let $\ell(h_1), \ell(h_2)$ be it's respective arclength. The Fréchet distance is defined as,

$$d_F(h_1, h_2) = \inf\{\sup|h_1(\sigma_1) - h_2(\sigma_2)|; \sigma_1, \sigma_2 \in \mathrm{Homeo}(S^1)\} + |\ell(h_1) - \ell(h_2)|.$$

For a given compact $W \subset \mathbb{C}$, the space of closed rectifiable curves $\mathscr{R}(S^1, W)$ is a complete metric space with respect to d_F.

Proposition 2.7 ([1])**.** *For a given compact $W \subset \mathbb{C}$, the space of closed rectifiable curves $\mathscr{R}(S^1, W)$ is complete metric space with respect to d_F.*

Proposition 2.8. *There exists a non-classical Schottky group Γ of $\mathfrak{D}_\Gamma = H_c$, and sequence of $\{\Gamma_n\}$ classical Schottky groups such that $\mathfrak{D}_{\Gamma_n} < \mathfrak{D}_\Gamma$ with $\sup \mathfrak{D}_{\Gamma_n} = \mathfrak{D}_\Gamma$.*

Proof. It follows from that the Hausdorff dimension map $\mathfrak{D} : \mathfrak{J}_g \to [0, 2]$ is real analytic map, and by the definition of H_c that, there exists a non-classical Γ with $\mathfrak{D}_\Gamma = H_c$. Theorem 1.4 implies $\mathfrak{J}_{g,o}^{H_c}$ is open submanifold of $\mathfrak{J}_g$, hence there exists a sequence of classical Schottkys $\{\Gamma_n\} \subset \mathfrak{J}_{o,g}^{H_c}$ with $\mathfrak{D}_{\Gamma_n} \to \mathfrak{D}_\Gamma$. □

3 Deformation Space and $\Psi(\Gamma_c)$

Take $\{\Gamma_n\}$ and Γ as given by Proposition 2.8. In particular, $\{\Gamma_n\}$ is a sequence of classical Schottky groups such that $\Gamma_n \to \Gamma$. Denote $QC(\mathbb{C})$ space of quasiconformal maps on $\mathbb{C}$. It follows from quasiconformal deformation theory of Schottky space, for $\Gamma_c \in \mathfrak{J}_{g,o}^{H_c}$ we can write ,

$$\mathfrak{J}_g \hat{=} \{f \circ \Gamma_c \circ f^{-1} \in \mathrm{PSL}(2, \mathbb{C}) | \quad \Gamma_c \quad \text{a classical Schottky}, f \in QC(\mathbb{C})\},$$

where "$\hat{=}$" denotes equality up to conjugacy by $\mathrm{PSL}(2, \mathbb{C})$.

Remark From *now on* throughout rest of the paper, we fix Γ_c to be a classical Schottky group a classical Schottky group with Hausdorff dimension $\mathfrak{D}_{\Gamma_c} < H_c$.

Hence there exists a sequence of quasiconformal maps f_n and f of $\mathbb{C}$ such that, $\Gamma_n = f_n(\Gamma)$ and $\Gamma = f(\Gamma_c)$. Here we write $f(\Gamma) := \{f \circ g \circ f^{-1} | g \in \Gamma\}$ for a given Kleinian group Γ and quasiconformal map f.

Proposition 3.1. *Let $\{\Gamma_n\} \subset \mathfrak{J}_g$ be a sequence of classical Schottky groups $\Gamma_n = f_n(\Gamma_c)$ such that $f_n \to f$, and set $\Gamma = f(\Gamma_c)$. Suppose for all n there exists a set classical generators S_n for Γ_n such that $\sup_{\gamma \in \cup S_n} |\operatorname{tr}(\gamma)| < \infty$. Then there exists a set of classical generators S of Γ_c such that for a subsequence $f_{i_n}(S)$ are classical generators of Γ_{i_n} and $\Gamma =< f(S) >$.*

Proof. For a given generating set S we use superscript notation S^n, to denote a sequence of distinct generating sets which are given by a sequence of Nielsen transformations of S. Recall that Nielsen transformations are automorphism of the free group.

Assume there exists no such set of classical generators S of Γ_c. Since $\Gamma_n, n \geq 1$ are classical Schottky groups, then any sequence $\{S_n\}$ of set of classical generators of Γ_n, for $\{S_n\}$ there exists $\{S^n\}$ a sequence of generating sets of Γ_c, which is not necessarily classical generators of Γ_c, such that, $S_n = f_n(S^n)$.

By assumption, we have S^n are distinct for large n. Since Schottky groups can be given as quasi-isometric embedding of Cayley graph of the free group on g generators with word metric into hyperbolic 3-space, implies that there exists $\gamma_n \in S^n$ such that, $|\operatorname{tr}(\gamma_n)| \to \infty$. Since $f_n \to f$, we must have $|\operatorname{tr}(f_n \circ \gamma_n \circ f_n^{-1}))| \to \infty$. This holds for all sets of classical classical generators of Γ_n, which is a contradiction, hence implies our result. □

Lemma 3.2. *Suppose $f_n \to f$ and $\Gamma = f(\Gamma_c)$. Every rectifiable quasi-circle of $\Gamma_n = f_n(\Gamma_c)$ is in $\mathscr{R}(S^1, W)$ for some compact W.*

Proof. Let η_n be a quasi-circle of Γ_n. Note $\Lambda_{\Gamma_n} \subset \eta_n$. Since limit set $\Lambda_{\Gamma_n} \to \Lambda_\Gamma$, and limit set $\Lambda(\Gamma)$ is compact, and η_n is rectifiable, we can find some compact set $W \supset \cup_n \Lambda_{\Gamma_n} \cup \Lambda_\Gamma$. □

Let E_n denote the collection of all *rectifiable* quasi-circles of $f_n(\Gamma_c)$. We define $\Psi(\Gamma_c) = \overline{\cup_n E_n}^{d_F}$ to be the closure of the set of all rectifiable quasi-circles of $f_n(\Gamma_c)$ in $\mathscr{R}(S^1, W)$.

Lemma 3.3. *Let $\Gamma = f(\Gamma_c)$ and $f_n \to f$. If η_Γ is a rectifiable quasi-circle of Γ then, $\eta_\Gamma \in \Psi(\Gamma_c)$.*

Proof. Since $\Gamma(\eta_\Gamma) = \eta_\Gamma$ and $\Gamma_n \to \Gamma$, we have $\inf_{\xi_n \in \Gamma_n(\eta_\Gamma)} \|\xi_n - \eta_\Gamma\|_{d_F} \to 0$. Set $\eta_n = f_n \circ f^{-1}(\eta_\Gamma)$. We have η_n is a quasi-circle of Γ_n. Since $f_n \circ f^{-1} \to id$ the identity map, and η_Γ is rectifiable which implies $\mathfrak{D}_\Gamma < 1$, hence we have that there exists some subsequence such that for n sufficiently large, $\mathfrak{D}_{\Gamma_n} < 1$.

This implies η_n rectifiable and $\|\eta_n - \eta_\Gamma\|_{d_F} \to 0$ for some sufficiently large subsequence. $\square$

We will denote by η_∞ a curve which is the limit of rectifiable quasi-circles of $f_n(\Gamma_c)$ in $\mathscr{R}(S^1, W)$.

Proposition 3.4. *Every η_∞ of $\mathscr{R}(S^1, W)$ is a quasi-circle of Γ. In addition, if Γ_n is a classical Schottky group for all n then, there exists a sequence $\{\eta_n^*\}$ of linear quasi-circles of Γ_n and, $\eta_\infty^* = \lim_n \eta_n^*$ is a quasi-circle of Γ.*

Proof. Since $\Gamma_n \to \Gamma$, it is obvious that η_∞ is a quasi-circle of Γ. Now assume that $\{\Gamma_n\}$ is a sequence of classical Schottky groups. To construct linear quasi-circles of Γ_n we proceed as follows:

Let S_n be a classical generating set of Γ_n. For $C_g \in \mathscr{C}_{S_n}^*$ and $\{c_i, c_{i+g}\} \in C_g$, take $p_{i,n} \in c_i, q_{i,n} \in c_{i+g}$ such that, $q_{i,n} \neq \gamma_{i,n}(p_{i,n})$, for $\gamma_{i,n} \in S_n, 1 \leq i \leq g$. Denote $p'_{i,n} = \gamma_{i,n}(p_{i,n})$. Let p_n, q_n denote the collection $\{p_{i,n}\}, \{q_{i,n}\}$ respectively.

For every $1 \leq i \leq g$, take $a_{i,p_{i,n},q_{i,n}}$ to be a finite piece-wise circular arc which connects $p_{i,n}$ to $q_{i,n}$, such that $a_{i,p_{i,n},q_{i,n}} \cap D_i = \emptyset$. In addition, let b_{i,q_n,p'_n} denote clock-wise arc along c_{i+g} connecting $p_{i,n}$ to $p'_{i,n}$. Define $a^*_{i,p_{i,n},q_{i,n}} = a_{i,p_{i,n},q_{i,n}} \cup b_{i,q_{i,n},p'_{i,n}}$. Since Γ_n is classical Schottky group and Mobius transformation of circular arc into circular arc, we have $a^*_{p_n,q_n} = \{a^*_{i,p_{i,n},q_{i,n}}\}$ is collection of disjoint curves consists of piece-wise circular arcs.

By construction, $\eta_n^j(a^*_{p_n,q_n})$ is piece-wise circular. Hence $\eta_n(a^*_{p_n,q_n})$ is linear quasi-circle of Γ_n. $\square$

Note that there are many linear quasi-circles of Γ_n. The importance of the particular type of linear quasi-circles η_n^* of Γ_n constructed in Proposition 3.4 is that, η_n^* depends on that Γ_n been classical Schottky group. This is crucial in the proof of our next result.

Lemma 3.5. *Let $\{\Gamma_n\}$ be a sequence of classical Schottky groups $\{f_n(\Gamma_c)\}$ with $\mathfrak{D}_{\Gamma_n} < H_c$. Let $\Gamma = f(\Gamma_c)$ be a non-classical Schottky group and $f_n \to f$. If $H_c < 1$, then there exists η_∞ that is not a quasi-circle of $f(\Gamma_c)$. In fact, we can take η_∞^* given by Proposition 3.4.*

Proof. Let S be any generating set of Γ. We denote by S^j the generating set obtained as $T_j(S)$ for a sequence of Nielsen transformations T_j. We will choose a sequence of T_j later in the proof.

Let $\eta_n(a^*_{p_n,q_n})$ be a linear quasi-circles with respect to a marking $C_{g,n} \in \mathscr{C}^*_{S_n}$ given by Proposition 3.4 for each n. We want to show that this sequence can't converge to a quasi-circle of Γ. Assume otherwise, then there exists a quasi-circle η_Γ of Γ such that for $\epsilon > 0$ and N_ϵ such that,

$$\|\eta_\Gamma - \eta_n(a^*_{p_n,q_n})\|_{d_F} < \epsilon, \quad \text{for} \quad n > N_\epsilon.$$

In particular, there exists J_ϵ with $\lim_{\epsilon\to 0} J_\epsilon \to \infty$ such that, η^j_Γ must be linear for $j < J_\epsilon$. Note that this dose not mean η_Γ itself is necessarily linear.

Let $C^j_g \in \mathscr{C}_{S^j}$. Since η^j_Γ consists of circular arcs for $j < J_\epsilon$, we claim that some arcs must lie on every $c_i \in C^j_g$:

In the following we parametrize all curves by $[0, 1]$. To prove the claim let's assume otherwise. Suppose for some $C_g \in \mathscr{C}_{S^j}$, there exists some $c_i, c'_i \in C_g$ such that $c_i \cap \eta_\Gamma = \{\alpha_k\}_1^{l_i}, c'_i \cap \eta_\Gamma = \{\alpha'_k\}_1^{l'_i}$ consists of l_i, l'_i points, and let $\gamma_i \in \Gamma$ with $\gamma_i(c_i) = c'_i$. Consider the curve $\eta^{j_i}_\Gamma$ at α_k. $\eta^{j_i}_\Gamma$ consists of finite circular arcs, and let $\rho_{i,o}$ be the piecewise circular arc of $\eta^{j_i}_\Gamma$ such that $\rho_{i,o} \cap D_i = \emptyset$. And let $\rho_{i,+}, \rho_{i,-}$ be the circular arc which is contained in D_i, D'_i and intersect with $\rho_{i,o}$ at α_k, α'_k. Set $\rho_i = \rho_{i,o} \cup \rho_{i,+} \cup \rho_{i,-}$.

There exists $\lambda_{i_n,n} \in \eta^{i_n}_n(a^*_{p_n,q_n})$ for some i_n and $n > N_\epsilon$ such that,

$$\sup\{|\lambda_{i_n,n}(\sigma_1) - \rho_i(\sigma_2)|\} < \epsilon, \quad \text{for some} \quad \sigma_1, \sigma_2 \in \mathrm{Aut}[0, 1].$$

Since $\lambda_{i_n,n}$ consist of finite circular arcs, by above inequality we must have that consist distinct circular arcs which intersect at points p_{1,i_n}, p_{2,i_n} close to α_k, α'_k. For $n > N_\epsilon$, let $\lambda_{o,i_n}, \lambda_{i_n,-}, \lambda_{i_n,+}$ denotes the circular arcs of $\lambda_{i_n,n}$ with intersection points $\lambda_{o,i_n} \cap \lambda_{i_n,-} = p_{1,i_n}, \lambda_{o,i_n} \cap \lambda_{i_n,+} = p_{2,i_n}$. By the construction of $\eta^{i_n}_n(a^*_{p_n,q_n})$, we can find $C_{g,i_n} \in \mathscr{C}^*_{S_n}$ and generating circles $c_{i_n,+}, c_{i_n,-} \in C_{g,i_n}$, such that there exists $\gamma_{i_n} \in \Gamma_n$ with $\gamma_{i_n}(c_{i_n,-}) = c_{i_n,+}$, and $\lambda_{i_n,+}, \lambda_{i_n,-}$ lie on $c_{i_n,+}, c_{i_n,-}$ respectively. This implies that $\gamma_{i_n}(\lambda_{i_n,-}) = \lambda_{i_n,+}$. Since $\rho_{i,+} \subset D_i, \rho_{i,-} \subset D'_i$ we must have, $c_{i_n,+} \cap D_i \neq \emptyset, c_{i_n,-} \cap D'_i \neq \emptyset$ for $n > N_\epsilon$. In fact, we have $\ell(c_{i_n,+} \cap D_i) + \epsilon \geq \ell(\rho_{i,+}), \ell(c_{i_n,-} \cap D'_i) + \epsilon \geq \ell(\rho_{i,-})$ for $n > N_\epsilon$, and ℓ denotes the arc-length.

On the other hand, since $\gamma_{i_n} \to \gamma_i$ and $\gamma_i(D_i) \cap D'_i = \emptyset$, hence we must have that, $\ell(c_{i_n,+} \cap D_i), \ell(c_{i_n,-} \cap D'_i) \to 0$ for $n \to \infty$, and since $\epsilon > 0$ is arbitrary we have $\ell(\rho_{i,-}), \ell(\rho_{i,+})$ are 0. This gives contradiction, which implies our claim.

For $\gamma \in \Gamma$ denote the complex rotational angle of γ by $\tau(\gamma)$. Next we note that there exists some generating set S' of Γ must contain a element with non-trivial complex rotation, a loxodromic transformation. To see this,

we simply recall that if not then, Γ is Fuchsian Schottky group which by [3], is classical Schottky group. In fact, the existence of S' implies there is a sequence of generating sets $\{S^j\}$ such that each S^j contains a loxodromic element. We choose T_j such that $T_j(S) = S^j$.

Let γ_j be a element of S^j which is of non-trivial, $\tau(\gamma_j) \neq 0$ complex rotation. Since Γ is non-classical Schottky group, $\psi(C_g^j) > 0$ for some $C_g^j \in \mathscr{C}_{S^j}$. This implies we have a pair of $c_{\gamma_j,+}, c_{\gamma_j,-} \in C_g^j$ with $\gamma_j(c_{\gamma_j,-}) = c_{\gamma_j,+}$ and $\gamma_j(D_{j,-}) \cap D_{j,+} = \emptyset$, for $\gamma_j \in S^j$. We have $\sigma_{\gamma_j} > 0$. This implies that we have a sequence $\{\gamma_j\}$ all of non-trivial complex rotation $\tau(\gamma_j) \bmod 2\pi \neq 0$ such that, for $c_{\gamma_j,\pm} \in C_g^j$ and $C_g^j \in \mathscr{C}_{S^j}$ with $\sigma_{\gamma_j} > 0$. By our first claim, we must have some arcs of η_Γ lie on $c_{\gamma_j,\pm}$. Denote these arcs by $b_{j,\pm}$. By $\tau(\gamma_j) \bmod 2\pi \neq 0$, there exists some j' so that, $b_{j',\pm}$ overlaps the non-circular part of $c_{\gamma_{j'},\pm}$. Let $b_{j'} \subset \eta_\Gamma^{j'}$ be this arc, the arc that lie on $c_{\gamma_{j'},\pm}$ consist of non-circular section. Hence there exists $\delta_{j'} > 0$ such that,

$$\sup\{|\lambda_{j'_n,n}(\sigma_1) - b_{j'}(\sigma_2)|\} > \delta_{j'}, \quad \text{for all} \quad \sigma_1, \sigma_2 \in \mathrm{Aut}[0,1], n \geq 0.$$

This implies that $\|\eta_\Gamma - \eta_n(a^*_{p_n,q_n})\|_{d_F} > \delta_{j'}, n \geq 0$. Therefore our result follows. □

4 Proof of Theorem 1.1

Proof. (Theorem 1.1)

We proof by contradiction. Suppose that $H_c < 1$.

First note that by Selberg Lemma we can just assume Kleinian group to be torsion-free.

Now note that if a Kleinian group Γ of $\mathfrak{D}_\Gamma < 1$ then it must be free. To show this, assume otherwise. Since $\mathfrak{D}_\Gamma < 1$ and Γ is purely loxodromic, it is convex-cocompact of second-kind. There exists a imbedded surface $\mathscr{R} = \Omega_\Gamma/\Gamma$ in $\mathbb{H}^3/\Gamma$. If $\mathscr{R}$ is incompressible then, subgroup $\pi_1(\mathscr{R}) \subset \Gamma$ have $\mathfrak{D}_{\pi_1(\mathscr{R})} = 1$ which is contradiction. If $\mathscr{R}$ is compressible then, we can cut along compression disks. We either end with incompressible surface as before or after finite many steps of cutting we obtain topological ball, which implies $\mathbb{H}^3/\Gamma$ is handle-body, hence free.

It follows from Proposition 2.8 and Proposition 3.1 there exists a sequence of classical Schottky groups $\Gamma_n =< f_n(S) >$ and a non-classical Schottky

group $\Gamma =< f(S) >$ such that, $f_n \to f$. By Lemma 3.2 and Proposition 3.4, every η_∞ of $\mathscr{R}(S^1, W)$ is quasi-circle of Γ.

Since $\eta^*_\infty \in \mathscr{R}(S^1, W)$ and by the fact that Γ is a non-classical Schottky group we have from Lemma 3.5, η^*_∞ can not be a quasi-circle of Γ. This gives our contradiction.

Finally, sharpness comes from the fact that, there exists Kleinian groups which is not free of Hausdorff dimension equal to one. Hence we have our result. □

E-mail: yonghou@princeton.edu

References

[1] Bott, R. (1968). Battelle Rencontres. Lectures in mathematics and physics. Eds. C, DeWitte and J, Wheeler. Benjamin, New York.

[2] Bowen, R. (1979). Hausdorff dimension of quasi-circles. I.H.E.S. 50. p. 11-25.

[3] Button, J. All Fuchsian Schottky groups are classical Schottky groups. Geometry&Topology Mono. Vol 1. p. 117-125.

[4] He, Y. M. Basmajian-type identities and Hausdorff dimension of limit sets. http://arxiv.org/abs/1602.06351

[5] Hou, Y. (2016). On smooth moduli space of Riemann surfaces.

[6] Hou, Y. (2010). Kleinian groups of small Hausdorff dimension are classical Schottky groups. I. Geometry & Topology. 14. p 473-519

[7] Hou, Y. (2014). All finitely generated Kleinian groups of small Hausdorff dimension are classical Schottky groups. http://arxiv.org/abs/1307.2677

[8] Schober, G. (1968). Continuity of curve functional and a technique involving quasiconformal mapping. Arch. Rat. Mech. Vol 29. pp. 378-389.